

9 Dunwoody Park, Suite 121 Dunwoody, GA 30338 P: 678-736-6374 F: 770-674-1701 Email: info@precisionpointdiagnostics.com www.precisionpointdiagnostics.com

## PATIENT INFO

NAME: Patient Sample REQUISITION ID: DPA213230010

# **PRECISION POINT** P88-Dietary Antigen Test

A Targeted Approach to Wellness



## **CLINIC INFO**

Sample Clinic ADDRESS: 121 Sample Lane Sample City, SS 10101

PHONE: (678)736-6374 FAX: (770)674-1701

# SUMMARY | 1/2

			ALLERGY				SENSIT	VITY	
DIETARY ANTIGEN	IgE	lgE Percent	lgG4	lgG4 Percent	IMMUNE TOLERANCE TO IgE	IgG	lgG Percent	C3d	C3d Percent
Almond	LOW	45%		2%			3%	HIGH	98%
Apple	LOW	47%		0%		LOW	14%		0%
Asparagus		9%		2%			6%	LOW	17%
Aspergillus Mix		0%		0%		LOW	15%	LOW	21%
Avocado		0%		0%			0%		0%
Banana	LOW	35%	LOW	18%		LOW	55%	LOW	27%
Barley	LOW	56%	LOW	39%	YES		3%		0%
Beef	LOW	30%		6%			0%	LOW	37%
Black Pepper	LOW	13%	LOW	15%	YES	LOW	15%		3%
Blueberry		0%	LOW	16%			10%		0%
Brewer's Yeast		0%		0%		LOW	55%		0%
Broccoli		1%	LOW	12%	YES	LOW	48%		8%
Cabbage		0%	LOW	28%			0%	LOW	44%
Сасао	LOW	29%		0%		LOW	19%		0%
Candida	LOW	51%		0%		LOW	27%		1%
Cantaloupe		0%		0%	YES		0%		0%
Carrot	LOW	14%	LOW	13%	YES		0%		8%
Casein	LOW	35%	LOW	34%	YES	LOW	52%		0%
Cashew	LOW	24%		1%			0%	MODERATE	82%
Cauliflower		0%	LOW	71%			0%		0%
Celery		0%		0%			0%		0%
Cherry		0%	MODERATE	80%	YES	LOW	15%		0%
Chicken		0%		2%			0%		0%
Cinnamon		0%		0%			9%		0%
Clam	HIGH	>99%		3%		LOW	12%	MODERATE	89%
Coconut	LOW	41%		0%			1%	MODERATE	81%
Codfish		0%	HIGH	>99%	YES	LOW	20%		8%
Coffee		0%	LOW	26%	YES	LOW	40%	LOW	13%
Corn	LOW	33%		0%			0%		3%
Cottonseed		0%	LOW	38%			1%		0%
Cow's Milk	LOW	63%	LOW	29%	YES	LOW	18%	LOW	26%
Crab		0%		1%			0%		0%
Cucumber		0%		0%			0%		0%
Egg Albumin	HIGH	93%		8%			1%	LOW	32%
Egg Yolk		0%	LOW	43%	YES	LOW	12%	LOW	13%
English Walnut		0%	LOW	56%		LOW	25%	LOW	34%
Flax Seed		0%	LOW	53%			2%		0%
Flounder		0%	LOW	58%		LOW	32%		0%

# SUMMARY | 2/2

			ALLERGY				SENSITI	/ITY	
DIETARY ANTIGEN	IgE	lgE Percent	lgG4	IgG4 Percent	IMMUNE TOLERANCE TO IgE	IgG	lgG Percent	C3d	C3d Percent
Garlic		0%	MODERATE	76%			6%		5%
Ginger		0%	LOW	42%	YES	LOW	13%		5%
Gluten	HIGH	>99%		0%			1%	LOW	67%
Goat's Milk	MODERATE	84%	LOW	52%		LOW	49%	LOW	58%
Grapefruit		0%		0%	YES		0%		0%
Grapes		0%	LOW	70%	YES	LOW	13%		0%
Green Olive		0%	LOW	57%	YES		1%		0%
Green Pea		0%		4%	YES		4%		0%
Green Pepper		0%	LOW	25%			0%		0%
Halibut		0%	LOW	50%			3%		0%
Honeydew		0%		0%		LOW	48%		0%
Hops		0%		0%			1%		0%
Kidney Bean	LOW	25%	LOW	26%	YES	LOW	11%	LOW	27%
Lemon		0%		0%			0%		0%
Lettuce	LOW	37%	LOW	18%	YES		0%		0%
Lima Bean	LOW	20%	LOW	16%	YES		0%	LOW	27%
Lobster	LOW	69%		0%			0%		0%
Mushroom	LOW	16%		0%			0%		0%
Mustard	LOW	46%	LOW	23%			1%		0%
Navy Bean	LOW	73%	LOW	59%	YES		8%	LOW	24%
Oat	LOW	14%	2011	0%	120		9%	2011	0%
Onion	2011	8%		0%			0%		0%
Orange	LOW	14%	LOW	14%	YES		3%		0%
Peach	LOW	0%	LOW	0%			0%		0%
Peanut		0%		1%	YES		2%		0%
Pear		0%		0%			0%		0%
Pecan		0%	LOW	60%		LOW	15%		0%
Pineapple		0%	LOW	0%		LOW	0%		0%
Plum	LOW	29%		0%			0%		0%
Pork	LOW	0%	LOW	72%		LOW	13%	LOW	19%
Rice		0%	LOW	0%		LOW	5%	LOW	0%
Rye	LOW	31%		0%			5%		0%
Salmon	LOW	0%	HIGH	98%			0%		0%
Scallops	HIGH	95%	mon	0%			0%		0%
Sesame	nion	0%		0%		LOW	46%		0%
Shrimp		4%		0%		LOW	0%	LOW	42%
Soybean		0%		0%	YES		0%	HIGH	97%
Spinach	LOW	12%	LOW	21%	YES		1%	LOW	15%
Strawberry	LOW	0%	LOW	0%	TES		0%	LOW	0%
String Bean		0%	LOW	32%			0%		0%
Sweet Potato		0%	LOW	31%			0%		0%
			LOW						
Tea Tomato		0% 0%		0% 0%		LOW	26% 0%		0% 0%
	MODERATE	94%	MODERATE		YES		1%		0%
Tuna	WIUDERATE		WODERATE	89%	TES				
Turkey		0%		8%			0%		0%
Vanilla		0%		0%			6%		0%
Watermelon		0%	1014	0%	┟────┤		0%		0%
White Potato		0%	LOW	50%	┟────┤		0%		3%
Whole Wheat		0%		9%			0%		0%
Yellow Squash		0%	MODERATE	89%			0%	LOW	20%

## LESS RESTRICTIVE DIET

The Less Restrictive Diet removes foods with high levels of reactivity for IgE and IgG. The Less Restrictive Diet rotates foo ds with moderate IgG reactivity where levels of C3d are also present due to increased inflammatory potential.

High IgG4 foods are listed separately, as IgG4 is not generally inflammatory, and its role is largely favorable apart from a handful of conditions. This allows the provider to determine whether to remove these foods based on the individual patient. The red "Remove at Provi ders Discretion" column reflects only IgG4 immunogenicity. Refer to "Understanding The P88 Dietary Antigen Test Results" guide for an expanded list of conditions associated with IgG4-RDs.

NO LIMI	TATION	ROTATE	ELIMINATE	ELIMINATE (IgG4	
These foods produce no immune reaction within your system at this time		These foods should be rotated out of your diet for a period of 72 hrs or reduced in overall intake.	Remove these foods entirely from your diet.	Remove at Provider's Discretion	
Almond	Hops		Barley	Codfish	
Apple	Kidney Bean		Clam	Salmon	
Asparagus	Lemon		Egg Albumin		
Aspergillus Mix	Lettuce		Gluten		
Avocado	Lima Bean		Rye		
Banana	Lobster		Scallops		
Beef	Mushroom		Whole Wheat		
Black Pepper	Mustard				
Blueberry	Navy Bean				
Brewer's Yeast	Oat				
Broccoli	Onion				
Cabbage	Orange				
Cacao	Peach				
Candida	Peanut				
Cantaloupe	Pear				
Carrot	Pecan				
Casein	Pineapple				
Cashew	Plum				
Cauliflower	Pork				
Celery	Rice				
Cherry	Sesame				
Chicken	Shrimp				
Cinnamon	Soybean				
Coconut	Spinach				
Coffee	Strawberry				
Corn	String Bean				
Cottonseed	Sweet Potato				
Cow's Milk	Теа				
Crab	Tomato				
Cucumber	Tuna				
Egg Yolk	Turkey				
English Walnut	Vanilla				
Flax Seed	Watermelon				
Flounder	White Potato				
Garlic	Yellow Squash				
Ginger					
Goat's Milk					
Grapefruit					
Grapes					
Green Olive					
Green Pea					
Green Pepper					
Halibut					
Honeydew				1	

This test has been developed and its performance characteristics determined by Precision Point Diagnostics. It has not been cleared by the U.S. Food and Drug Administration.

Lab Director: Steven Lobel, PhD

## **MORE RESTRICTIVE DIET**

The More Restrictive Diet removes foods with high and moderate levels of IgE, IgG, and complement (C3d). Additionally, low IgG reactivity with any positive complement response are rotated because C3d has the potential to amplify an IgG reaction 1000-10,000-fold.

High and moderate IgG4 foods are listed separately, as IgG4 is not generally inflammatory, and its role is largely favorable apart from a handful of conditions. This allows the provider to determine whether to remove these foods based on the individual patient. The red "Remove at Providers Discretion" column reflects only IgG4 immunogenicity. Refer to "Understanding The P88 Dietary Antigen Test Results" guide for an expanded list of conditions associated with IgG4-RDs.

NO LIMI	TATION	ROTATE	ELIMINATE	ELIMINATE (IgG4)		
These foods produce within your syste		These foods should be rotated out of your diet for a period of 72 hrs or reduced in overall intake.	Remove these foods entirely from your diet.	Remove at Provider's Discretion		
Apple	Peanut	Aspergillus Mix	Almond	Cherry		
Asparagus	Pear	Banana	Barley	Codfish		
Avocado	Pecan	Coffee	Cashew	Garlic		
Beef	Pineapple	Cow's Milk	Clam	Salmon		
Black Pepper	Plum	Egg Yolk	Coconut	Yellow Squash		
Blueberry	Rice	English Walnut	Egg Albumin			
Brewer's Yeast	Sesame	Kidney Bean	Gluten			
Broccoli	Shrimp	Pork	Goat's Milk			
Cabbage	Spinach		Rye			
Cacao	Strawberry		Scallops			
Candida	String Bean		Soybean			
Cantaloupe	Sweet Potato		Tuna			
Carrot	Теа		Whole Wheat			
Casein	Tomato					
Cauliflower	Turkey					
Celery	Vanilla					
Chicken	Watermelon					
Cinnamon	White Potato					
Corn						
Cottonseed						
Crab						
Cucumber						
Flax Seed						
Flounder						
Ginger						
Grapefruit						
Grapes						
Green Olive						
Green Pea						
Green Pepper						
Halibut						
Honeydew						
Hops						
Lemon Lettuce						
Lima Bean						
Lobster						
Mushroom						
Mustard						
Navy Bean						
Oat						
Onion						
Orange						
Peach						

This test has been developed and its performance characteristics determined by Precision Point Diagnostics. It has not been cleared by the U.S. Food and Drug Administration.

Lab Director: Steven Lobel, PhD

**Patient Sample** 

DPA213230010 DRAFT DATE:

## IMMUNE INDEX

The Precision 88 is the only dietary antigen test to categorize overall reactivity of foods by adjusting for immunogenicity across four independent markers: IgE, IgG4, total IgG, and C3d (complement). Our immunogenicity-adjusted algorithm, known here as the Immune Index, emphasizes C3d, and de-emphasizes IgG4. This specialized calculation generates its own rank of most-to-least reactive foods and allows for consideration of increased flexibility towards IgG4 reactive foods in the absence of IgG4-RDs.

Concurrently, the red "Remove at Providers Discretion" columns on pp. 3 and 4 reflect only IgG4 immunogenicity. Refer to pp. 4-5 in our Understanding The P88 Dietary Antigen Test Results guide, for an expanded list of conditions associated with IgG4-RDs.

Denk	DIETARY	Immune
Rank	ANTIGEN	Index
1	Clam	MODERATE
2	Goat's Milk	MODERATE
3	Almond	LOW
4	Banana	LOW
5	Egg Albumin	LOW
6	Gluten	LOW
7	Kidney Bean	LOW
8	Cow's Milk	LOW
9	Cashew	LOW
10	Coconut	LOW
11	Apple	LOW
12	Aspergillus Mix	LOW
13	Beef	LOW
14	Black Pepper	LOW
15	Cacao	LOW
16	Candida	LOW
17	Coffee	LOW
18	Egg Yolk	LOW
19	English Walnut	LOW
20	Navy Bean	LOW
21	Lima Bean	LOW
22	Pork	LOW
23	Scallops	LOW
24	Soybean	LOW
25	Spinach	LOW
26	Casein	LOW
27	Codfish	LOW
28	Tuna	LOW
29	Asparagus	
30	Barley	
31	Brewer's Yeast	
32	Broccoli	
33	Cabbage	
34	Carrot	
35	Cherry	
36	Corn	
37	Flounder	
38	Grapes	
39	Honeydew	
40	Lettuce	
41	Ginger	
42	Lobster	
43	Mushroom	
44	Mustard	

DIETARY	Immune
ANTIGEN	Index
Orange	
Yellow Squash	
Salmon	
Avocado	
Blueberry	
Cantaloupe	
Celery	
Chicken	
Cinnamon	
Cottonseed	
Crab	
Cucumber	
Cauliflower	
Garlic	
Grapefruit	
Green Olive	
Green Pea	
Green Pepper	
Halibut	
Flax Seed	
Lemon	
Onion	
Hops	
Peach	
Rice	
Strawberry	
Tomato	
Vanilla	
Watermelon	
	ANTIGEN Oat Orange Pecan Plum Rye Sesame Shrimp Yellow Squash Tea Salmon Avocado Blueberry Cantaloupe Celery Chicken Cinnamon Cottonseed Crab Cucumber Cauliflower Garlic Grapefruit Green Pepper Halibut Flax Seed Lemon Onion Hops Peach Peanut Pear Pineapple Rice Strawberry String Bean Sweet Potato Tomato Turkey Vanilla

This test has been developed and its performance characteristics determined by Precision Point Diagnostics. It has not been cleared by the U.S. Food and Drug Administration. Lab Director: Steven Lobel, PhD GA Clinical License: 044-160 5

#### **BIOGENIC COMPOUNDS**

This table recognizes the dynamics of symptom-eliciting compounds as potential, non-immune-response-driven, explanations for perturbances, irritations and allergy-mimicking reactions. Reactive foods that also populate for these compounds can identify additional patterns of food reactions that are not mediated by IgE or IgG. For example, several reactions in a category may signal an intolerance not previously considered, or may confirm observed symptomologies and metabolic disturbances, thus prompting a dietary source review for those and similar-acting compounds. This illustration reminds of the myriad of reasons why biological systems respond to food (and other environmental) triggers.

DIETARY ANTIGEN	Oxalates	Amines	Glutamate	Histamine	Lectins	Nitrite	FOD-MAP	Phenol	Salicylates
Almond		Н							Н
Apple									
Asparagus									
Avocado									
Banana									
Barley									
Blueberry									
Broccoli									
Cabbage									
Casein									
Cashew							М		
Cauliflower									
Celery									
Coconut						М			
Coffee									
Corn									
Grapefruit									
Kidney Bean									
Lettuce									
Mushroom									
Navy Bean									
Onion									
Orange									
Peach									
Peanut									
Pear									
Pineapple									
Plum									
Shrimp									
Soybean	Н			Н			Н		
Spinach									
Strawberry									
Теа									
Tomato									
Turkey									
Watermelon									
White Potato									
Whole Wheat									
whole wheat									





9 Dunwoody Park, Suite 121 Dunwoody, GA 30338 P: 678-736-6374 F: 770-674-1701 Email: info@precisionpointdiagnostics.com www.precisionpointdiagnostics.com

# **P88-Dietary Antigen Test**

A Targeted Approach to Wellness

## PATIENT INFO

NAME: Patient Sample REQUISITION ID: DPA213230010

## **CLINIC INFO**

Sample Clinic ADDRESS: 121 Sample Lane Sample City, SS 10101

PHONE: (678)736-6374 FAX: (770)674-1701

IgE/IgG4 Food Allergies

## **Dietary Antigen Exposure by Food Group**

	IgE	lgG4	
Meats, Dairy	LOW	LOW	Meats, Dairy
Poultry	MODERATE	LOW	Vegetables Poultry HIGH
Legumes, Beans	LOW	LOW	Fruits MODERATE Legumes, Beans
Fungi	LOW	LOW	
Fish	LOW	MODERATE	
Shellfish	MODERATE	LOW	Seeds, Nuts Fungi
Herbs, Spices,	LOW	LOW	
Grains, Grasses	LOW	LOW	Grains, Grasses Fish
Seeds, Nuts	LOW	LOW	Herbs, Spices, Shellfish Seasonings
Fruits	LOW	LOW	Immune
Vegetables	LOW	LOW	Tolerance to IgE

## **Dietary Antigen Exposure by Food Group**

In this test, a human serum sample is probed for the presence of IgE and IgG4 antibodies which have an exact affinity for specific dietary allergens. Dietary allergens are clustered by the food groups shown in the table and graph above. The quantitative summation of the IgE and IgG4 results within the offending food groups are expressed graphically. The exclusion of the offending food group(s) from the diet has been shown to reduce the severity of symptoms associated with food allergies.

### **Immune Tolerance To IgE**

In high levels, IgG4 antibodies alone can trigger an immune response within the body. However, data is available that provides support for the notion that IgG4 can serve another specific function of controlling antigen recognition by IgE and consequently regulating anaphylactic reactions and IgE-mediated immunity. IgG4 can act as a blocking agent by preventing IgE from binding to targeted receptor sites and releasing histamine. We refer to this as the Immune Tolerance to IgE.

IgE/IgG4 Food Allergies

## **Understanding the Key**

It is important to understand how reactive your patient is to a given food. Based on peer-reviewed literature and the methodology used in our test, the lower 10% of reactivity is likely asymptomatic and represents the reference range of a normal/negative result in the general population. The HIGH range represents the top 5% of reactivity, and MODERATE the next 20%. Thus, the HIGH and MODERATE ranges combined represent the top 25% of reactivity. A LOW result represents the range of reactivity between 10% and 75% of the population.

Some foods have a greater prevalence of reactivity in the general population, most notably, dairy and casein, wheat and glute n, shellfish, tree nuts, and eggs. The increased prevalence of allergies and sensitivities to these foods is reflected in our test as an adjust ment of the HIGH range to the top 10% of the sample population, the MODERATE range the next 40%, and a LOW result represents the range between 10% and 50% of the population.

#### lgE

The IgE antibody response is the most commonly known food allergy response. This response usually occurs immediately and can create severe symptoms such as swelling, hives, itching, and - in some cases - anaphylaxis. Even though IgE reactions are immediate, the allergic potential of food-based allergens can remain in your system 1-2 days after ingestion, extending the presence of symptoms during this duration. IgE reactions can be permanent or they may improve with the elimination diet and gut treatment. IgE reactions stimulate the release of histamine in the body.

#### lgG4

IgG4, which is a subclass of IgG, is a distinct antibody in the immune system. IgG4 total antibody is important in relation to IgE because this antibody acts as a blocking agent for an IgE reaction. When the IgG4 reaction is greater than the IgE reaction for a particular antigen, IgG4 blocks the IgE antibodies from binding to the receptor sites and releasing histamine, thereby reducing severity of the sympto ms associated with the IgE reaction. This is referred to as the blocking potential. IgG4 carries its own clinical relevance in high level s and may mediate several conditions and diseases.

ANTIGEN	RESULT	lgE (μg/mL)	REF. RANGE	IMMUNE TOLERANCE TO IgE	ANTIGEN	RESULT	lgG4 (μg/mL)	REF. RANGE	
	1	<b>NEATS, DA</b>	IRY		MEATS, DAIRY				
Beef	1.50	LOW	<0.54 µg/ml		Beef	0.37		<0.76 µg/ml	
Casein	0.55	LOW	<0.29 µg/ml	YES	Casein	2.36	LOW	<0.56 µg/ml	
Cow's Milk	2.18	LOW	<0.3 µg/ml	YES	Cow's Milk	3.16	LOW	<0.6 µg/ml	
Goat's Milk	2.12	MODERATE	<0.25 µg/ml		Goat's Milk	1.74	LOW	<0.25 µg/ml	
Pork	0.00		<0.43 µg/ml		Pork	3.06	LOW	<0.36 µg/ml	
		POULTRY	Y			POUL	TRY		
Chicken	0.00		<0.39 µg/ml		Chicken	0.33		<0.64 µg/ml	
Egg Albumin	24.05	HIGH	<3.01 µg/ml		Egg Albumin	6.31		<6.88 µg/ml	
Egg Yolk	0.09		<0.24 µg/ml	YES	Egg Yolk	3.83	LOW	<0.87 µg/ml	
Turkey	0.00		<0.26 µg/ml		Turkey	0.27		<0.39 µg/ml	
	LE	GUMES, BI	EANS		LEGUMES, BEANS				
Green Pea	0.07		<0.32 µg/ml	YES	Green Pea	0.20		<0.32 µg/ml	
Kidney Bean	0.20	LOW	<0.15 µg/ml	YES	Kidney Bean	0.75	LOW	<0.34 µg/ml	
Lima Bean	0.38	LOW	<0.25 µg/ml	YES	Lima Bean	0.42	LOW	<0.35 µg/ml	
Navy Bean	2.89	LOW	<0.97 µg/ml	YES	Navy Bean	3.14	LOW	<0.8 µg/ml	
Peanut	0.11		<0.86 µg/ml	YES	Peanut	0.59		<1.54 µg/ml	
Soybean	0.10		<1.65 µg/ml	YES	Soybean	0.51		<2.04 µg/ml	
String Bean	0.00		<0.22 µg/ml		String Bean	1.75	LOW	<0.63 µg/ml	
		FUNGI				FUN	GI		
Aspergillus Mix	0.06		<0.27 µg/ml		Aspergillus Mix	0.00		<0.56 µg/ml	
Brewer's Yeast	0.00		<0.28 µg/ml		Brewer's Yeast	0.00		<0.36 µg/ml	
Candida	1.60	LOW	<0.61 µg/ml		Candida	0.00		<0.33 µg/ml	
Mushroom	0.32	LOW	<0.25 µg/ml		Mushroom	0.00		<0.55 µg/ml	
		FISH				FIS	H		
Codfish	0.09		<0.22 µg/ml	YES	Codfish	8.19	HIGH	<0.34 µg/ml	
Flounder	0.00		<0.29 µg/ml		Flounder	1.44	LOW	<0.37 µg/ml	
Halibut	0.00		<0.27 µg/ml		Halibut	1.28	LOW	<0.31 µg/ml	
Salmon	0.00		<0.27 µg/ml		Salmon	4.68	HIGH	<0.25 µg/ml	
Tuna	2.07	MODERATE	<0.28 µg/ml	YES	Tuna	2.33	MODERATE	<0.21 µg/ml	

## **Patient Results**

IgE/IgG4 Food Allergies

# **Patient Results**

ANTIGEN	RESULT	lgE (μg/mL)	REF. RANGE	IMMUNE TOLERANCE TO IgE	ANTIGEN	RESULT	lgG4 (μg/mL)	REF. RANGE
		SHELLFIS	H			SHELL		
Clam	19.52	HIGH	<3.14 µg/ml		Clam	1.03		<1.86 µg/ml
Crab	0.00		<0.4 µg/ml		Crab	0.17		<0.54 µg/ml
Lobster	1.14	LOW	<0.19 µg/ml		Lobster	0.00		<0.27 µg/ml
Scallops	2.76	HIGH	<0.47 µg/ml		Scallops	0.00		<0.31 µg/ml
Shrimp	0.12		<0.15 µg/ml		Shrimp	0.00		<0.28 μg/ml
ennip	-	SPICES, SE	ASONINGS				SEASONIN	
Black Pepper	0.27	LOW	<0.21 µg/ml	YES	Black Pepper	0.34	LOW	<0.32 µg/ml
Cinnamon	0.00		<0.14 µg/ml		Cinnamon	0.00		<0.42 µg/ml
Garlic	0.00		<0.24 µg/ml		Garlic	2.47	MODERATE	<0.36 µg/ml
Ginger	0.04		<0.26 µg/ml	YES	Ginger	2.25	LOW	<0.39 µg/ml
Hops	0.03		<0.25 µg/ml		Hops	0.00		<0.48 µg/ml
Mustard	0.79	LOW	<0.35 µg/ml		Mustard	0.65	LOW	<0.35 µg/ml
Vanilla	0.00		<0.2 µg/ml		Vanilla	0.00		<0.29 µg/ml
	GF	RAINS, GRA	ASSES	1		GRAINS, G	GRASSES	1.0
Barley	0.52	LOW	<0.18 µg/ml	YES	Barley	0.59	LOW	<0.23 µg/ml
Corn	0.55	LOW	<0.26 µg/ml		Corn	0.09		<0.44 µg/ml
Gluten	18.38	HIGH	<3.47 µg/ml		Gluten	0.21		<6.18 µg/ml
Oat	0.26	LOW	<0.21 µg/ml		Oat	0.00		<0.27 µg/ml
Rice	0.00		<0.19 µg/ml		Rice	0.10		<0.28 µg/ml
Rye	0.48	LOW	<0.27 µg/ml		Rye	0.00		<0.44 µg/ml
Whole Wheat	0.00		<0.32 µg/ml		Whole Wheat	0.40		<0.42 µg/ml
		SEEDS, NU				SEEDS,	NUTS	PO/
Almond	1.13	LOW	<0.27 µg/ml		Almond	0.25		<0.53 µg/ml
Сасао	0.42	LOW	<0.2 μg/ml		Cacao	0.00		<0.34 µg/ml
Cashew	0.57	LOW	<0.36 µg/ml		Cashew	0.12		<0.51 µg/ml
Coffee	0.10	2011	<0.32 µg/ml	YES	Coffee	0.44	LOW	<0.24 µg/ml
Cottonseed	0.00		<0.19 µg/ml		Cottonseed	0.80	LOW	<0.29 μg/ml
English Walnut	0.00		<0.21 µg/ml		English Walnut	1.56	LOW	<0.26 µg/ml
Flax Seed	0.00		<0.49 µg/ml		Flax Seed	1.79	LOW	<0.47 µg/ml
Pecan	0.00		<0.39 μg/ml		Pecan	1.47	LOW	<0.36 µg/ml
Sesame	0.00		<0.15 µg/ml		Sesame	0.00	2011	<0.24 µg/ml
		FRUITS	p.8,			FRU	TS	PO/
Apple	0.59	LOW	<0.23 µg/ml		Apple	0.03		<0.25 µg/ml
Avocado	0.00		<0.38 µg/ml		Avocado	0.00		<0.54 µg/ml
Banana	0.43	LOW	<0.21 µg/ml		Banana	0.38	LOW	<0.33 µg/ml
Blueberry	0.00		<0.33 µg/ml		Blueberry	0.71	LOW	<0.54 µg/ml
Cantaloupe	0.00		<0.28 µg/ml	YES	Cantaloupe	0.01		<0.32 µg/ml
Cherry	0.03		<0.35 µg/ml	YES	Cherry	2.19	MODERATE	<0.33 µg/ml
Coconut	0.82	LOW	<0.32 µg/ml		Coconut	0.00		<0.46 µg/ml
Cucumber	0.00		<0.15 µg/ml		Cucumber	0.00		<0.23 µg/ml
Grapefruit	0.07		<0.15 µg/ml	YES	Grapefruit	0.14		<0.29 µg/ml
Grapes	0.07		<0.15 µg/ml	YES	Grapes	0.96	LOW	<0.23 µg/ml
Green Olive	0.05		<0.2 µg/ml	YES	Green Olive	1.28	LOW	<0.29 µg/ml
Green Pepper	0.00		<0.19 µg/ml		Green Pepper	0.43	LOW	<0.24 µg/ml
Honeydew	0.00		<0.22 μg/ml		Honeydew	0.00	2011	<0.38 µg/ml
Lemon	0.00		<0.15 µg/ml		Lemon	0.00		<0.19 µg/ml
Orange	0.22	LOW	<0.19 µg/ml	YES	Orange	0.37	LOW	<0.32 µg/ml
Peach	0.00		<0.29 μg/ml		Peach	0.00		<0.22 µg/ml
Pear	0.00	<u> </u>	<0.18 μg/ml		Pear	0.00	1	<0.22 μg/ml
Pineapple	0.00	1	<0.16 µg/ml		Pineapple	0.00		<0.19 μg/ml
Plum	0.00	LOW	<0.19 µg/ml		Plum	0.00	ł	<0.19 µg/ml
Strawberry	0.00	1000	<0.19 μg/ml		Strawberry	0.00	ł	<0.24 μg/ml
Tomato	0.00		<0.28 μg/ml		Tomato	0.00	ł	<0.33 µg/mi <0.21 µg/ml
		<u> </u>					<u> </u>	
Watermelon	0.00		<0.25 µg/ml		Watermelon	0.00	MODERATE	<0.36 µg/ml
Yellow Squash	0.00		<0.22 µg/ml		Yellow Squash	2.42	MODERATE	<0.32 µg/ml

IgE/IgG4 Food Allergies

# **Patient Results**

ANTIGEN	RESULT	lgE (µg/mL)	REF. RANGE	IMMUNE TOLERANCE TO IgE	ANTIGEN	RESULT	lgG4 (µg/mL)	REF. RANGE
		VEGETABI	LES			VEGET	ABLES	
Asparagus	0.31		<0.32 µg/ml		Asparagus	0.13		<0.36 µg/ml
Broccoli	0.11		<0.32 µg/ml	YES	Broccoli	0.66	LOW	<0.53 µg/ml
Cabbage	0.00		<0.18 µg/ml		Cabbage	0.41	LOW	<0.25 µg/ml
Carrot	0.23	LOW	<0.19 µg/ml	YES	Carrot	0.26	LOW	<0.24 µg/ml
Cauliflower	0.00		<0.16 µg/ml		Cauliflower	2.36	LOW	<0.32 µg/ml
Celery	0.00		<0.25 µg/ml		Celery	0.03		<0.3 µg/ml
Lettuce	0.39	LOW	<0.17 µg/ml	YES	Lettuce	0.41	LOW	<0.32 µg/ml
Onion	0.13		<0.15 µg/ml		Onion	0.00		<0.23 µg/ml
Spinach	0.22	LOW	<0.22 µg/ml	YES	Spinach	0.71	LOW	<0.47 µg/ml
Sweet Potato	0.00	1	<0.33 µg/ml		Sweet Potato	0.81	LOW	<0.37 µg/ml
Теа	0.00		<0.15 µg/ml		Теа	0.00		<0.23 µg/ml
White Potato	0.00		<0.22 µg/ml		White Potato	1.56	LOW	<0.36 µg/ml



9 Dunwoody Park, Suite 121 Dunwoody, GA 30338 P: 678-736-6374 F: 770-674-1701 Email: info@precisionpointdiagnostics.com www.precisionpointdiagnostics.com

### **PATIENT INFO**

NAME: Patient Sample REQUISITION ID: DPA213230010

# PRECISION POINT P88-Dietary Antigen Test

A Targeted Approach to Wellness

**CLINIC INFO** 


Sample Clinic ADDRESS: 121 Sample Lane Sample City, SS 10101

PHONE: (678)736-6374 FAX: (770)674-1701

IgG/C3d Food Sensitivities

## **Dietary Antigen Exposure by Food Group**

	lgG
Meats, Dairy	LOW
Poultry	LOW
Legumes, Beans	LOW
Fungi	LOW
Fish	LOW
Shellfish	LOW
Herbs, Spices,	LOW
Grains, Grasses	LOW
Seeds, Nuts	LOW
Fruits	LOW
Vegetables	LOW



### Dietary Antigen Exposure by Food Group

In this test, a human serum sample is probed for the presence of IgG antibodies which have an exact affinity for specific dietary allergens. Dietary allergens are clustered by the food groups shown in the table and graph above. The quantitative summation of the IgG results within the offending food groups are expressed graphically. The exclusion of the offending food group(s) from the diet has been shown to reduce the severity of symptoms associated with food allergies.

## IgG/C3d Food Sensitivities

## **Understanding the Key**

It is important to understand how reactive your patient is to a given food. Based on peer-reviewed literature and the methodology used in our test, the lower 10% of reactivity is likely asymptomatic and represents the reference range of a normal/negative result in the general population. The HIGH range represents the top 5% of reactivity, and MODERATE the next 20%. Thus, the HIGH and MODERATE ranges combined represent the top 25% of reactivity. A LOW result represents the range of reactivity between 10% and 75% of the population.

Some foods have a greater prevalence of reactivity in the general population, most notably, dairy and casein, wheat and gluten, shellfish, tree nuts, and eggs. The increased prevalence of allergies and sensitivities to these foods is reflected in our test as an adjustment of the HIGH range to the top 10% of the sample population, the MODERATE range the next 40%, and a LOW result represents the range between 10% and 50% of the population.

#### lgG

The IgG antibody response creates sensitivity to a particular food. Symptoms may be less severe than with IgE allergic reaction and can manifest anywhere from 3-72 hours after exposure. IgG reactions create inflammation that makes many pathologies worse. The delayed response makes sensitivities difficult to identify without a diagnostic test. Sensitivities can improve with treatment and improved gut health.

### <u>C3d</u>

C3d is a complement antigen and an activator of our complement cascade system. Reaction to the specified food will worsen if C3d activation is present along with an IgG antibody response. The C3 protein attaches to the antigen and amplifies the IgG response, increasing the inflammatory potential of IgG titer. Complement is not dependent on exposurevi or antibody presence, and represents innate immune function.

ANTIGEN	RESULT	lgG (μg/mL)	REF. RANGE	ANTIGEN	RESULT	C3d (µg/mL)	REF. RANGE	
	MEA	TS, DAIRY			MEA	TS, DAIRY		
Beef	0.00		<2.59 µg/ml	Beef	0.62	LOW	<0.22 µg/ml	
Casein	30.70	LOW	<2.93 µg/ml	Casein	0.07		<0.23 µg/ml	
Cow's Milk	38.34	LOW	<15.86 µg/ml	Cow's Milk	0.68	LOW	<0.33 µg/ml	
Goat's Milk	16.34	LOW	<2.91 µg/ml	Goat's Milk	0.79	LOW	<0.17 µg/ml	
Pork	3.90	LOW	<2.36 µg/ml	Pork	0.36	LOW	<0.21 µg/ml	
	PC	DULTRY		POULTRY				
Chicken	0.00		<1.24 µg/ml	Chicken	0.02		<0.16 µg/ml	
Egg Albumin	3.78		<17.73 µg/ml	Egg Albumin	0.90	LOW	<0.42 µg/ml	
Egg Yolk	2.48	LOW	<2.17 µg/ml	Egg Yolk	0.79	LOW	<0.68 µg/ml	
Turkey	0.00		<0.84 µg/ml	Turkey	0.00		<0.19 µg/ml	
	LEGUN	/IES, BEANS		LEGUMES, BEANS				
Green Pea	0.81		<1.5 µg/ml	Green Pea	0.00		<0.24 µg/ml	
Kidney Bean	2.05	LOW	<1.92 µg/ml	Kidney Bean	0.19	LOW	<0.12 µg/ml	
Lima Bean	0.00		<2.1 µg/ml	Lima Bean	0.30	LOW	<0.18 µg/ml	
Navy Bean	2.99		<4.38 µg/ml	Navy Bean	0.24	LOW	<0.15 µg/ml	
Peanut	1.37		<3.7 µg/ml	Peanut	0.00		<0.33 µg/ml	
Soybean	0.00		<2.7 µg/ml	Soybean	3.31	HIGH	<0.58 µg/ml	
String Bean	0.18		<3.03 µg/ml	String Bean	0.00		<0.18 µg/ml	
FUNGI				FUNGI				
Aspergillus Mix	32.10	LOW	<23.71 µg/ml	Aspergillus Mix	0.40	LOW	<0.27 µg/ml	
Brewer's Yeast	26.56	LOW	<3.97 µg/ml	Brewer's Yeast	0.00		<0.14 µg/ml	
Candida	57.31	LOW	<17.42 µg/ml	Candida	0.12		<0.16 µg/ml	
Mushroom	1.28		<15.73 µg/ml	Mushroom	0.33		<1.29 µg/ml	
		FISH		FISH				
Codfish	1.52	LOW	<0.97 µg/ml	Codfish	0.22		<0.26 µg/ml	
Flounder	2.03	LOW	<0.73 µg/ml	Flounder	0.00		<0.16 µg/ml	
Halibut	0.15		<0.52 µg/ml	Halibut	0.00		<0.21 µg/ml	
Salmon	0.00		<0.8 µg/ml	Salmon	0.00		<0.15 µg/ml	
Tuna	0.27		<0.76 µg/ml	Tuna	0.00		<0.12 µg/ml	

## **Patient Results**

# IgG/C3d Food Sensitivities

# **Patient Results**

ANTIGEN	RESULT	lgG (μg/mL)	REF. RANGE	ANTIGEN	RESULT	C3d (µg/mL)	REF. RANGE
	SHI	ELLFISH			SH	ELLFISH	
Clam	10.35	LOW	<8.28 µg/ml	Clam	1.45	MODERATE	<0.24 µg/ml
Crab	0.00		<1.38 µg/ml	Crab	0.00		<0.14 µg/ml
Lobster	0.00		<1.42 µg/ml	Lobster	0.00		<0.16 µg/ml
Scallops	0.00		<0.96 µg/ml	Scallops	0.00		<0.14 µg/ml
Shrimp	0.00		<1.28 µg/ml	Shrimp	0.23	LOW	<0.13 µg/ml
	HERBS, SPIC	ES, SEASONIN			HERBS, SPIC	ES, SEASONIN	
Black Pepper	15.3	LOW	<11.4 µg/ml	Black Pepper	0.1		<0.15 µg/ml
Cinnamon	3.0		<3.21 µg/ml	Cinnamon	0.0		<0.15 µg/ml
Garlic	0.4		<1.2 µg/ml	Garlic	0.1		<0.15 µg/ml
Ginger	13.8	LOW	<12.06 µg/ml	Ginger	0.2		<0.33 µg/ml
Hops	0.2	-	<1.89 µg/ml	Hops	0.0		<0.23 µg/ml
Mustard	0.2		<1.38 μg/ml	Mustard	0.0		<0.18 µg/ml
Vanilla	6.8		<9.54 µg/ml	Vanilla	0.0		<0.15 µg/ml
Vanna		S, GRASSES	<0.0 4 μ <sub>6</sub> / m	Vanna		S, GRASSES	(0.10 µ6/m
Barley	0.24		<0.77 µg/ml	Barley	0.05	5, GIA3515	<0.14 µg/ml
Corn	0.24		<1.81 µg/ml	Corn	0.05		<0.14 µg/ml
Gluten	8.03		<54.14 μg/ml	Gluten	0.12	LOW	<0.19 µg/ml
Oat	0.75		<0.81 µg/ml	Oat	0.00	LOW	<0.12 μg/ml
Rice	0.75		<1.13 μg/ml	Rice	0.10		<0.12 µg/ml
	1.40		<1.13 μg/ml	Rye	0.10		<0.13 μg/ml
Rye Whole Wheat	-			Whole Wheat			
whole wheat	0.00		<1.39 µg/ml	whole wheat	0.02		<0.14 µg/ml
A lue e u el		OS, NUTS	11 56	A lue e us el		DS, NUTS	10.24
Almond	0.49	1014	<1.56 µg/ml	Almond	1.95	HIGH	<0.24 µg/ml
Cacao	16.48	LOW	<9.31 µg/ml	Cacao	0.05		<0.12 µg/ml
Cashew	0.00		<2.1 µg/ml	Cashew	0.85	MODERATE	<0.14 µg/ml
Coffee	20.88	LOW	<6.72 μg/ml	Coffee	0.33	LOW	<0.29 µg/ml
Cottonseed	0.64		<3.19 µg/ml	Cottonseed	0.05		<0.18 µg/ml
English Walnut	6.66	LOW	<1.88 µg/ml	English Walnut	1.14	LOW	<0.49 µg/ml
Flax Seed	0.75		<2.31 µg/ml	Flax Seed	0.00		<0.16 µg/ml
Pecan	1.52	LOW	<1.11 µg/ml	Pecan	0.00		<0.14 µg/ml
Sesame	2.25	LOW	<0.3 µg/ml	Sesame	0.00		<0.12 µg/ml
		RUITS				RUITS	
Apple	0.55	LOW	<0.46 µg/ml	Apple	0.05		<0.14 µg/ml
Avocado	0.98		<3.13 µg/ml	Avocado	0.07		<0.63 µg/ml
Banana	4.16	LOW	<0.79 µg/ml	Banana	0.20	LOW	<0.15 µg/ml
Blueberry	1.94		<1.98 µg/ml	Blueberry	0.07		<0.19 µg/ml
Cantaloupe	0.04		<1.18 µg/ml	Cantaloupe	0.02		<0.22 µg/ml
Cherry	0.92	LOW	<0.64 µg/ml	Cherry	0.05		<0.18 µg/ml
Coconut	0.35		<2.25 µg/ml	Coconut	0.58	MODERATE	<0.14 µg/ml
Cucumber	0.00		<0.38 µg/ml	Cucumber	0.06		<0.13 µg/ml
Grapefruit	0.10		<1.15 µg/ml	Grapefruit	0.03		<0.14 µg/ml
Grapes	0.53	LOW	<0.49 µg/ml	Grapes	0.00		<0.12 µg/ml
Green Olive	0.24		<1.93 µg/ml	Green Olive	0.00		<0.14 µg/ml
Green Pepper	0.00		<0.45 µg/ml	Green Pepper	0.00		<0.15 µg/ml
Honeydew	2.50	LOW	<0.51 µg/ml	Honeydew	0.00		<0.2 µg/ml
Lemon	0.00		<0.19 µg/ml	Lemon	0.09		<0.12 µg/ml
Orange	0.44		<1.34 µg/ml	Orange	0.00		<0.12 µg/ml
Peach	0.00		<0.75 µg/ml	Peach	0.00		<0.14 µg/ml
Pear	0.00		<0.45 µg/ml	Pear	0.00		<0.16 µg/ml
Pineapple	0.00		<0.33 µg/ml	Pineapple	0.00		<0.12 µg/ml
Plum	0.00		<0.78 µg/ml	Plum	0.00		<0.12 µg/ml
Strawberry	0.04		<0.88 μg/ml	Strawberry	0.00		<0.18 µg/ml
Tomato	0.00		<0.27 μg/ml	Tomato	0.00		<0.13 µg/ml
Watermelon	0.00		<0.27 μg/ml	Watermelon	0.03		<0.23 µg/ml

## IgG/C3d Food Sensitivities

# **Patient Results**

ANTIGEN	RESULT	lgG (μg/mL)	REF. RANGE	ANTIGEN	RESULT	C3d (µg/mL)	REF. RANGE
VEGETABLES				VEGETABLES			
Asparagus	4.18		<7.25 µg/ml	Asparagus	0.24	LOW	<0.19 µg/ml
Broccoli	5.41	LOW	<1.73 µg/ml	Broccoli	0.13		<0.14 µg/ml
Cabbage	0.00		<0.37 µg/ml	Cabbage	0.29		<0.13 µg/ml
Carrot	0.21		<1.12 µg/ml	Carrot	0.13		<0.14 µg/ml
Cauliflower	0.00		<0.78 µg/ml	Cauliflower	0.00		<0.14 µg/ml
Celery	0.00		<1.72 µg/ml	Celery	0.00		<0.17 µg/ml
Lettuce	0.12		<0.83 µg/ml	Lettuce	0.00		<0.13 µg/ml
Onion	0.00		<0.2 µg/ml	Onion	0.00		<0.12 µg/ml
Spinach	0.21		<1.21 µg/ml	Spinach	0.27	LOW	<0.2 µg/ml
Sweet Potato	0.24		<1.94 µg/ml	Sweet Potato	0.10		<0.42 µg/ml
Теа	4.70	LOW	<1.92 µg/ml	Теа	0.00		<0.13 µg/ml
White Potato	0.66		<3.69 µg/ml	White Potato	0.41		<0.68 µg/ml

